

SURFACE LINKS WHICH ARE COVERINGS OVER THE STANDARD TORUS

INASA NAKAMURA

ABSTRACT. We consider surface links in the 4-space or the 4-sphere which can be described in braid forms over the standard torus, which we call torus-covering-links. Torus-covering-links include spun T^2 -knots, turned spun T^2 -knots, symmetry-spun tori and torus T^2 -knots. In this paper we give examples of interesting torus-covering-links, namely, torus-covering T^2 -links which are ribbon but not symmetry-spun T^2 -links, and a torus-covering T^2 -knot whose triple point number is positive.

0. INTRODUCTION

Closed 2-manifolds embedded locally flatly in the Euclidean 4-space $\mathbb{R}^4$ or the 4-sphere S^4 are called *surface links*. It is known that any oriented surface link can be described in a form of the closure of a simple surface braid, i.e. in a braid form over the standard 2-sphere (cf. [12, 14]).

As surface knots of genus one which can be made from classical knots, there are *spun T^2 -knots*, *turned spun T^2 -knots*, *symmetry-spun tori* and *torus T^2 -knots*. Consider $\mathbb{R}^4$ as obtained by rotating $\mathbb{R}_+^3$ around the boundary $\mathbb{R}^2$. Then a *spun T^2 -knot* is obtained by rotating a classical knot (cf. [3]), a *turned spun T^2 -knot* by turning a classical knot once while rotating it (cf. [3]), a *symmetry-spun torus* by turning a classical knot with periodicity rationally while rotating (cf. [20]), and a *torus T^2 -knot* is a surface knot on the boundary of a neighborhood of a solid torus in $\mathbb{R}^4$ (cf. [10]). Symmetry-spun tori include spun T^2 -knots, turned spun T^2 -knots and torus T^2 -knots. We call the link version of a symmetry-spun torus, a spun T^2 -knot, and a turned spun T^2 -knot a *symmetry-spun T^2 -link*, a *spun T^2 -link* and a *turned spun T^2 -link* respectively. We enumerate several properties of symmetry-spun T^2 -links.

- (0.1) A symmetry-spun T^2 -link is equivalent to a spun T^2 -link or a turned spun T^2 -link (cf. [20]).
- (0.2) A spun T^2 -link is ribbon and the turned spun T^2 -link of a non-trivial classical link is not ribbon (cf. [3, 19]).
- (0.3) A symmetry-spun T^2 -link is pseudo-ribbon and has the triple point number zero (cf. [15]).
- (0.4) A symmetry-spun T^2 -link has a classical link group (cf. [20]).

Now we consider oriented surface links which can be described in braid forms over the standard torus, which we will define as *torus-covering-links* (see Definition 2.1). By definition, a torus-covering-link is associated with a *torus-covering-chart*, which is a chart on the standard torus. Torus-covering-links include symmetry-spun T^2 -links (and spun T^2 -links, turned spun T^2 -links, and torus T^2 -links). A torus-covering-link has no 2-knot component. Each component of a torus-covering-link

1991 *Mathematics Subject Classification*. Primary 57Q45; Secondary 57Q35.

Key words and phrases. surface link; 2-dimensional braid; knot group; triple point number; quandle cocycle invariant.

is of genus at least one. There is a natural question.

(Q) What difference from symmetry-spun T^2 -links do torus-covering-links have?

In Section 2, we consider torus-covering-links in S^4 , and from Section 3 throughout this paper we consider torus-covering T^2 -links in $\mathbb{R}^4$, i.e. torus-covering-links in $\mathbb{R}^4$ whose each component is of genus one. Torus-covering T^2 -links are associated with torus-covering-charts without black vertices.

In Section 2 we define torus-covering-links (Definition 2.1), and *turned torus-covering-links* (Definition 2.4).

In Section 3 we study link groups of torus-covering T^2 -links. We show that “*There are infinitely many 2-component torus-covering T^2 -links whose link groups are not classical link groups.*” (Theorem 3.5). We show its knot version as well: “*There are infinitely many torus-covering T^2 -knots whose knot groups are not classical knot groups.*” (Theorem 3.9). This means that torus-covering T^2 -links are not always symmetry-spun T^2 -links. Concerning Theorem 3.5, we have a corollary: “*There are infinitely many 2-component torus-covering T^2 -links whose link groups are not 2-component 2-link groups.*” (Corollary 3.8).

In Section 4 we show that the torus-covering-links given in Theorems 3.5 and 3.9 are ribbon, i.e. “*There are infinitely many torus-covering T^2 -links which are ribbon but not symmetry-spun T^2 -links.*” (Theorem 4.1).

Symmetry-spun T^2 -links are pseudo-ribbon and by Theorem 4.1 the torus-covering T^2 -links given in Theorems 3.5 and 3.9 are ribbon. Both of them have the triple point number zero. In Section 5 we show that “*There is a torus-covering T^2 -knot whose triple point number is positive.*” (Theorem 5.1). Moreover we have “*There is a torus-covering T^2 -knot which is not (-)-amphicheiral.*” (Corollary 5.2).

The author would like to thank professors Takashi Tsuboi and Elmar Vogt for suggesting this problem, and professor Akio Kawauchi for advising this title.

1. DEFINITIONS AND PRELIMINARIES

Definition 1.1. A closed 2-manifold embedded locally flatly in $\mathbb{R}^4$ or S^4 is called a *surface link*. A surface link with one component is called a *surface knot*. A surface link whose each component is of genus zero (resp. one) is called a *2-link* (resp. *T^2 -link*). In particular a surface knot of genus zero (resp. one) is called a *2-knot* (resp. *T^2 -knot*).

An oriented surface link F is *trivial* (or *unknotted*) if there is an embedded 3-manifold M with $\partial M = F$ such that each component of M is a handlebody.

Let $\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a generic projection. Then a *surface diagram* of a surface link F in $\mathbb{R}^4$ is the image $\pi(F)$ with additional crossing information at the singularity set. An oriented surface link F is called *pseudo-ribbon* if there is a surface diagram of F whose singularity set consists of double points, and *ribbon* if F is obtained from a trivial 2-link F_0 by 1-handle surgeries along a finite number of mutually disjoint 1-handles attaching to F_0 . By definition, a ribbon surface link is pseudo-ribbon.

Two surface links are *equivalent* if there is an ambient isotopy of $\mathbb{R}^4$ (or S^4) or an orientation-preserving self-homeomorphism of $\mathbb{R}^4$ (or S^4) which deforms one to the other.

Definition 1.2. A compact and oriented 2-manifold S_B embedded properly and locally flatly in $D_1^2 \times D_2^2$ is called a *braided surface* of degree m if S_B satisfies the following conditions:

- (i) $\text{pr}_2|_{S_B} : S_B \rightarrow D_2^2$ is a branched covering map of degree m ,
- (ii) ∂S_B is a closed m -braid in $D_1^2 \times \partial D_2^2$, where D_1^2, D_2^2 are 2-disks, and $\text{pr}_2 : D_1^2 \times D_2^2 \rightarrow D_2^2$ is the projection to the second factor.




FIGURE 1.1. Vertices in a chart

A braided surface S_B is called a *surface braid* if ∂S_B is the trivial closed braid. Moreover, S_B is called *simple* if every singular index is two. Two braided surfaces of the same degree are *equivalent* if there is a fiber-preserving ambient isotopy of $D_1^2 \times D_2^2 \text{ rel } D_1^2 \times \partial D_2^2$ which carries one to the other.

There is a theorem which corresponds to Alexander's theorem for classical oriented links.

Theorem 1.3 ([12, 14]). *Any oriented surface link in $\mathbb{R}^4$ can be deformed by an ambient isotopy of $\mathbb{R}^4$ to the closure of a simple surface braid.*

There is a *chart* which represents a simple surface braid.

Definition 1.4. Let m be a positive integer, and Γ be a graph on a 2-disk D_2^2 . Then Γ is called a *surface link chart* of degree m if it satisfies the following conditions:

- (i) $\Gamma \cap \partial D_2^2 = \emptyset$.
- (ii) Every edge is oriented and labeled, and the label is in $\{1, \dots, m-1\}$.
- (iii) Every vertex has degree 1, 4, or 6.
- (iv) At each vertex of degree 6, there are six edges adhering to which, three consecutive arcs oriented inward and the other three outward, and those six edges are labeled i and $i+1$ alternately for some i .
- (v) At each vertex of degree 4, the diagonal edges have the same label and are oriented coherently, and the labels i and j of the diagonals satisfy $|i-j| > 1$ (Fig. 1.1).

A vertex of degree 1 (resp. 6) is called a *black vertex* (resp. *white vertex*). A black vertex (resp. white vertex) in a chart corresponds to a branch point (resp. triple point) in the surface diagram of the associated simple surface braid by the projection pr_2 .

A chart with a boundary represents a simple braided surface.

There is a notion of *C-move equivalence* between two charts of the same degree. The following theorem is well-known.

Theorem 1.5 ([13, 14]). *Two charts of the same degree are C-move equivalent if and only if their associated simple braided surfaces are equivalent.*

2. TORUS-COVERING-LINKS

In this section, we greatly rely on [20]. We define *torus-covering-links* (Definition 2.1) and *turned torus-covering-links* (Definition 2.4). Throughout this paper, let $\sigma_1, \sigma_2, \dots, \sigma_{m-1}$ be the standard generators of the braid group of degree m .

Definition 2.1. Let D^2 be a 2-disk and $S^1 = [0, 1]/\sim$ with $0 \sim 1$. First, embed $D^2 \times S^1 \times S^1$ into S^4 or $\mathbb{R}^4$ naturally, or more precisely, consider as follows (cf. [20], see also [3, 16]). Let $S^1 \times S^1$ be a standardly embedded torus in S^4 and let $D^2 \times S^1 \times S^1$ be a tubular neighborhood of $S^1 \times S^1$ in S^4 . We can assume that its framing is canonical, i.e. the homomorphism induced by the inclusion map




FIGURE 2.1. A torus-covering-link

$H_1(\{p\} \times S^1 \times S^1; \mathbb{Z}) \rightarrow H_1(S^4 - S^1 \times S^1; \mathbb{Z})$ where $p \in \partial D^2$, is zero. Let $\bar{l} = \partial D^2 \times \{0\} \times \{0\}$, $\bar{s} = \{p\} \times S^1 \times \{0\}$, and $\bar{r} = \{p\} \times \{0\} \times S^1$ be curves on $\partial D^2 \times S^1 \times S^1$. Put $E^4 = \text{cl}(S^4 - D^2 \times S^1 \times S^1)$. Let l, s and r be the canonical curves on ∂E^4 , which are identified with $\bar{l}, \bar{s}$ and $\bar{r}$ under the natural identification map $i : \partial D^2 \times S^1 \times S^1 \rightarrow \partial E^4$. The curves l, s and r represent a basis of $H_1(\partial E^4; \mathbb{Z})$. Let $f : \partial E^4 \rightarrow E^4$ be a homeomorphism such that $f_{\#} \begin{pmatrix} l & s & r \end{pmatrix} = \begin{pmatrix} l & s & r \end{pmatrix} A^f$, where $A^f \in GL(3, \mathbb{Z}) \cong \pi_0 \text{Homeo}(\partial E^4)$. The map f can be extended to a homeomorphism $\tilde{f} : E^4 \rightarrow E^4$ if and only if $A^f \in H$, where

$$H = \left\{ \begin{pmatrix} \pm 1 & 0 & 0 \\ * & \alpha & \gamma \\ * & \beta & \delta \end{pmatrix} \in GL(3, \mathbb{Z}); \alpha + \beta + \gamma + \delta \equiv 0 \pmod{2} \right\}.$$

Now consider $D^2 \times S^1 \times S^1$ to be embedded in $S^4 = E^4 \cup_i D^2 \times S^1 \times S^1$ or $\mathbb{R}^4 = (E^4 - \{*\}) \cup_i (D^2 \times S^1 \times S^1)$, where $*$ $\in \text{Int} E^4$, and identify $D^2 \times S^1 \times S^1$ with $D^2 \times I_3 \times I_4 / \sim$, where $(x, 0, v) \sim (x, 1, v)$ and $(x, u, 0) \sim (x, u, 1)$ for $x \in D^2, u \in I_3 = [0, 1]$ and $v \in I_4 = [0, 1]$.

Let us consider a surface link S embedded in $D^2 \times S^1 \times S^1$ such that $S \cap (D^2 \times I_3 \times I_4)$ is a simple braided surface. We will call S a *torus-covering-link* (Fig. 2.1). In particular, if each component of a torus-covering-link is of genus one, we call it a *torus-covering T^2 -link*.

A torus-covering-link is associated with a chart on the standard torus, i.e. with a chart Γ_T on $I_3 \times I_4$ with $\Gamma_T \cap (I_3 \times \{0\}) = \Gamma_T \cap (I_3 \times \{1\})$ and $\Gamma_T \cap (\{0\} \times I_4) = \Gamma_T \cap (\{1\} \times I_4)$. Denote the classical braids associated with $\Gamma_T \cap (I_3 \times \{0\})$ and $\Gamma_T \cap (\{0\} \times I_4)$ by a and b respectively. We will call Γ_T a *torus-covering-chart with boundary braids a and b* . In particular, a torus-covering T^2 -link is associated with a torus-covering-chart Γ_T without black vertices, and the torus-covering T^2 -link is determined from the boundary braids a and b , which are commutative. We will call Γ_T a *torus-covering-chart without black vertices and with boundary braids a and b* .

By definition, torus-covering-links include symmetry-spun T^2 -links.

As we stated in Theorem 1.5, for two surface link charts of the same degree, their associated surface links are equivalent if the charts are C-move equivalent. It follows that if two torus-covering-charts of the same degree are C-move equivalent, then their associated torus-covering-links are equivalent.

A torus-covering-link has no 2-knot component. Each component of a torus-covering-link is of genus at least one.

Let Γ_T be a torus-covering-chart of degree m with the trivial boundary braids. Let F be the surface link associated with the surface link chart obtained from Γ_T by assuming it to be a surface link chart. Then the torus-covering-link associated with the torus-covering-chart Γ_T is obtained from F by applying m trivial 1-handle surgeries.

Example 2.2. (2.2.1) Let Γ_T be the torus-covering-chart of degree 2 without black vertices and with boundary braids σ_1^3 and e (the trivial braid). Then the torus-covering-knot associated with Γ_T is the spun T^2 -knot of a right-handed trefoil.

(2.2.2) Let Γ_T be the torus-covering-chart of degree 2 without black vertices and with boundary braids σ_1^3 and σ_1^3 . Then the torus-covering-knot associated with Γ_T is the turned spun T^2 -knot of a right-handed trefoil.

(2.2.3) Let Γ_T be the torus-covering-chart of degree 2 without black vertices and with boundary braids σ_1^2 and σ_1^3 . Then the torus-covering-knot associated with Γ_T is a symmetry-spun torus.

Let Γ_T be a torus-covering-chart on $I_3 \times I_4$, where $I_3 = I_4 = [-1, 1]$. Consider a map $\underline{\rho} : I_3 \times I_4 \rightarrow I_3 \times I_4$ such that $\underline{\rho}(t_1, t_2) = (-t_2, t_1)$. We call $\underline{\rho}(\Gamma_T)$ the torus-covering-chart obtained from Γ_T by *rotating it by $\pi/2$* . Let us denote by ρ the $\pi/2$ rotation, which maps the torus-covering-link S associated with Γ_T to the torus-covering-link $\rho(S)$ associated with $\underline{\rho}(\Gamma_T)$.

Lemma 2.3 (cf. [20]). *Let S be the torus-covering-link associated with a torus-covering-chart Γ_T . Then the torus-covering-links $\rho^n(S)$ obtained from S by the $n\pi/2$ rotation for $n \in \mathbb{Z}$ are all equivalent.*

Proof. The torus-covering-links $\rho^n(S)$ are in $D^2 \times S^1 \times S^1$ in $S^4 = E^4 \cup_i D^2 \times S^1 \times S^1$, where $S^1 = [-1, 1]/\sim$ with $-1 \sim 1$. We will show that S and $\rho(S)$ are equivalent. Consider a homeomorphism $f : D^2 \times S^1 \times S^1 \rightarrow D^2 \times S^1 \times S^1$ such that $f(x, t_1, t_2) = (x, -t_2, t_1)$. The map $f|_{\partial D^2 \times S^1 \times S^1}$ can be considered as an

orientation-preserving self-homeomorphism of ∂E^4 . Since $A^f = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \in$

H , $f|_{\partial D^2 \times S^1 \times S^1}$ can be extended to $\tilde{f} : E^4 \rightarrow E^4$. Hence there is an orientation-preserving self-homeomorphism of S^4 which deforms S to $\rho(S)$. $\square$

Remark. Teragaito proved in [20] the same theorem in the symmetry-spun version. Lemma 7 in [20] corresponds to the $-\pi/2$ rotation.


We can consider *turned torus-covering-links*, which include turned spun T^2 -links (cf. [3, 20]).

Definition 2.4. We use the notations of Definition 2.1. Let $\sigma : \partial E^4 \rightarrow \partial E^4$ be the homeomorphism of a matrix $A^\sigma = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$. Then $E^4 \cup_{\sigma i} D^2 \times S^1 \times S^1$

is homeomorphic to S^4 . Let Γ_T be a torus-covering-chart and let S be the torus-covering-link associated with Γ_T in $S^4 = E^4 \cup_i D^2 \times S^1 \times S^1$. Then we can consider the torus-covering-link obtained from S by changing the identification map i to σi , which we will call the *turned torus-covering-link associated with (S, Γ_T) or S* , and use the notation $\tau(S, \Gamma_T)$ or $\tau(S)$. That is, we define $\tau(S)$ as follows:

$$(E^4 \cup_{\sigma i} D^2 \times S^1 \times S^1 \cong S^4, S) = (S^4 = E^4 \cup_i D^2 \times S^1 \times S^1, \tau(S)).$$

Note that for a spun T^2 -link S , $\tau(S)$ is the turned spun T^2 -link. Let us denote by τ the map which maps S to the turned torus-covering-link $\tau(S)$.

FIGURE 2.2. The torus-covering-charts Γ_T and $\Gamma_{\tau(T)} = \underline{\tau}(\Gamma_T)$

By definition, we have the following lemmas.

Lemma 2.5. *Let S be the torus-covering-link associated with a torus-covering-chart Γ_T , and consider $\tau(S)$, the turned torus-covering-link associated with S . Then the torus-covering-chart $\Gamma_{\tau(T)}$ associated with $\tau(S)$ is as in Fig. 2.2.*

Proof. Consider the inverse map $\sigma^{-1} : \partial E^4 \rightarrow \partial E^4$ of σ . Then $\sigma_{\#}^{-1} \begin{pmatrix} l & s & r \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} l & s & r \end{pmatrix}$. The homeomorphism σ^{-1} is decomposed as $1 \times \underline{\tau}$ for a homeomorphism $\underline{\tau} : S^1 \times S^1 \rightarrow S^1 \times S^1$ such that $\underline{\tau}_{\#} \begin{pmatrix} s & r \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} s & r \end{pmatrix}$. Hence, regarding torus-covering-charts as graphs on the standard torus, the torus-covering-chart $\Gamma_{\tau(T)}$ is $\underline{\tau}(\Gamma_T)$, which is as in Fig. 2.2. $\square$

Lemma 2.6 (cf. [3, 20]). *For a torus-covering-link S , $\tau^2(S) = S$.*

Proof. Since $A^{\sigma^2} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix} \in H$, there is an orientation-preserving self-homeomorphism of S^4 which deforms S to $\tau^2(S)$. $\square$

The equivalence class of $\tau(S)$ is not determined from the equivalence class of S . There is a pair of torus-covering-links S and S' with the associated torus-covering-charts Γ_T and Γ'_T such that S and S' are equivalent but $\tau(S, \Gamma_T)$ and $\tau(S', \Gamma'_T)$ are not equivalent (Proposition 2.7). In Proposition 2.7, let $S' = \rho(S)$. By Lemma 2.3, S and S' are equivalent. By Proposition 2.7, $\tau(S')$ (i.e. $\tau \circ \rho(S)$) and $\rho \circ \tau(S)$ are not equivalent. However, by Lemma 2.3, $\rho \circ \tau(S)$ and $\tau(S)$ are equivalent. Hence $\tau(S')$ and $\tau(S)$ are not equivalent.

Proposition 2.7. *There is a torus-covering-chart Γ_T such that $\tau \circ \rho(S)$ and $\rho \circ \tau(S)$ are not equivalent, where S is the torus-covering-knot associated with Γ_T , ρ is the $\pi/2$ rotation and τ is the map which maps a torus-covering-link to its associated turned torus-covering-link.*

Proof. Consider the torus-covering-chart Γ_T of degree 2 without black vertices and with boundary braids σ_1^3 and e . The torus-covering-knot S associated with Γ_T is the spun T^2 -knot of a trefoil. Then the boundary braids of the torus-covering-chart $\underline{\tau} \circ \underline{\rho}(\Gamma_T)$ are e and σ_1^3 , and by Lemma 2.3 $\tau \circ \rho(S)$ is equivalent to S . On the other hand, by Lemma 2.3 again, $\rho \circ \tau(S)$ is equivalent to $\tau(S)$, which is the turned spun T^2 -knot of the trefoil. By (0.2), $\tau \circ \rho(S)$ and $\rho \circ \tau(S)$ are not equivalent. $\square$

3. KNOT GROUPS AND LINK GROUPS

From now on throughout this paper, we consider torus-covering T^2 -links in $\mathbb{R}^4$, i.e. the torus-covering-links in $\mathbb{R}^4$ associated with torus-covering-charts without black vertices. We can compute link groups of torus-covering T^2 -links (Lemma 3.4).

Before stating Lemma 3.4, we will give the definition of *Artin's automorphism* (Definition 3.3, cf. [14]). Let D^2 be a 2-disk, β an m -braid in a cylinder $D^2 \times [0, 1]$, Q_m the starting point set of β .

Definition 3.1. An *isotopy of D^2 associated with β* is an ambient isotopy $\{\phi_t\}_{t \in [0, 1]}$ of D^2 such that

- (i) $\phi_t|_{\partial D^2} = \text{id}$,
- (ii) $\phi_t(Q_m) = \text{pr}_1(\beta \cap \text{pr}_2^{-1}(t))$ for $t \in [0, 1]$, where $\text{pr}_1 : D^2 \times [0, 1] \rightarrow D^2$ (resp. $\text{pr}_2 : D^2 \times [0, 1] \rightarrow [0, 1]$) is the projection to the first (resp. second) factor.

Definition 3.2. A *homeomorphism of D^2 associated with β* is the terminal map $\psi = \phi_1 : D^2 \rightarrow D^2$ of an isotopy $\{\phi_t\}_{t \in [0, 1]}$ of D^2 associated with β .

Let q_0 be a point of ∂D^2 . Identify the fundamental group $\pi_1(D^2 - Q_m, q_0)$ with the free group F_m generated by the standard generator system of $\pi_1(D^2 - Q_m, q_0)$.

Definition 3.3. *Artin's automorphism* of F_m associated with β is the automorphism of F_m induced by a homeomorphism of D^2 associated with β . We denote it by $\text{Artin}(\beta)$.

We can obtain Artin's automorphism (of the free group F_m associated with an m -braid) algebraically by the following rules. Let $F_m = \langle x_1, x_2, \dots, x_m \rangle$.

- (1) $\text{Artin}(\beta_1 \beta_2) = \text{Artin}(\beta_2) \circ \text{Artin}(\beta_1)$ for m -braids β_1 and β_2 , and
- (2)

$$\text{Artin}(\sigma_i)(x_j) = \begin{cases} x_j & \text{if } j \neq i, i+1, \\ x_i x_{i+1} x_i^{-1} & \text{if } j = i, \\ x_i & \text{if } j = i+1, \end{cases}$$

and

$$\text{Artin}(\sigma_i^{-1})(x_j) = \begin{cases} x_j & \text{if } j \neq i, i+1, \\ x_{i+1} & \text{if } j = i, \\ x_{i+1}^{-1} x_i x_{i+1} & \text{if } j = i+1, \end{cases}$$

where $i = 1, 2, \dots, m-1$ and $j = 1, 2, \dots, m$.

Lemma 3.4. Let Γ_T be a torus-covering-chart of degree m without black vertices and with boundary braids a and b . Let S be the torus-covering-link associated with Γ_T . Then the link group of S is obtained as follows:

$\pi_1(\mathbb{R}^4 - S) = \langle x_1, \dots, x_m \mid x_j = \text{Artin}(a)(x_j) = \text{Artin}(b)(x_j), j = 1, 2, \dots, m \rangle$, where $\text{Artin}(a) : F_m \rightarrow F_m$ (resp. $\text{Artin}(b)$) is Artin's automorphism of the free group $F_m = \langle x_1, \dots, x_m \rangle$ associated with the m -braid a (resp. b).

Proof. Let us denote by $D^2 \times S^1 \times S^1$ the neighborhood of the torus in which the torus-covering-link S is embedded as in Definition 2.1, where D^2 is a 2-disk and $S^1 = [0, 1]/\sim$ with $0 \sim 1$. Let $x_0 = \{q_0\} \times \{0\} \times \{0\}$ be the base point, where $q_0 \in \partial D^2$ and $\{0\} \in S^1$. Then $\pi_1(D^2 \times S^1 \times S^1 - S, x_0)$ is given by

$$(3.1) \quad \langle x_1, \dots, x_m, s, t \mid x_j = s \cdot \text{Artin}(a)(x_j) \cdot s^{-1}, x_j = t \cdot \text{Artin}(b)(x_j) \cdot t^{-1}, st = ts, j = 1, 2, \dots, m \rangle,$$

where s (resp. t) is represented by the meridian (resp. longitude) of $\{q_0\} \times S^1 \times S^1$.

Let B_0^3 be a 3-ball such that $(D^2 \times S^1) \cup B_0^3$ is a 3-ball and $(D^2 \times S^1) \cap B_0^3$ is homotopic to S^1 . Let us denote the 3-ball $(D^2 \times S^1) \cup B_0^3$ by E^3 . Put $X = E^3 \times S^1 - S$, $A = D^2 \times S^1 \times S^1 - S$ and $B = B_0^3 \times S^1$. We can assume that B and S are disjoint. Then $X = A \cup B$. We can also assume that the base point $x_0 = \{q_0\} \times \{0\} \times \{0\}$ is in $A \cap B$. Since A , B and $A \cap B$ are connected, by van Kampen's theorem, $\pi_1(X, x_0)$ is given by

$$[\pi_1(A, x_0) * \pi_1(B, x_0)] / \text{Vk}(X, A, B, x_0),$$

where $\text{Vk}(X, A, B, x_0)$ is the *van Kampen subgroup*, i.e. the smallest normal subgroup of the free product $\pi_1(A, x_0) * \pi_1(B, x_0)$ containing the elements of the form $i_*(\delta)j_*(\delta^{-1})$, where $i : A \cap B \rightarrow A$, $j : A \cap B \rightarrow B$ are inclusion maps, and $\delta \in \pi_1(A \cap B, x_0)$. Since $A \cap B$ is homotopic to $S^1 \times S^1$ and include $\{q_0\} \times S^1 \times S^1$, $\pi_1(A \cap B, x_0)$ has a presentation $\langle s'', t'' \mid s''t'' = t''s'' \rangle$ such that $i_*(s'') = s$ and $i_*(t'') = t$, where s and t are two of the generators of $\pi_1(A, x_0)$ with the presentation (3.1). On the other hand, $\pi_1(B, x_0)$ has the presentation $\langle t' \rangle$ such that $j_*(s'') = 1$ and $j_*(t'') = t'$. Hence $\pi_1(X) = \pi_1(E^3 \times S^1 - S)$ is

$$\langle x_1, \dots, x_m, t \mid x_j = \text{Artin}(a)(x_j), x_j = t \cdot \text{Artin}(b)(x_j) \cdot t^{-1}, j = 1, 2, \dots, m \rangle.$$

Since $E^3 \times S^1$ is naturally embedded in $\mathbb{R}^4$, the longitude of $\{q_0\} \times S^1 \times S^1$ which represents t bounds a disk in $\mathbb{R}^4$. Hence we have

$$\pi_1(\mathbb{R}^4 - S) = \langle x_1, \dots, x_m \mid x_j = \text{Artin}(a)(x_j) = \text{Artin}(b)(x_j), j = 1, 2, \dots, m \rangle.$$

□

Not every link group of torus-covering T^2 -links is a classical link group, i.e. there are torus-covering T^2 -links which are not symmetry-spun T^2 -links (cf. (0.4)).

Theorem 3.5. *Let $\Gamma_{T,n}$ be the torus-covering-chart of degree 4 without black vertices and with boundary braids $\sigma_1\sigma_3$ and Δ^{2n} , where $\Delta = \sigma_1\sigma_2\sigma_3\sigma_1\sigma_2\sigma_1$ (Garside's Δ), and n is a positive integer. Let S_n be the torus-covering-link associated with $\Gamma_{T,n}$. Then the link group of S_n is not a classical link group. Moreover, S_n and S_m are not equivalent for $n \neq m$.*

Remark. The torus-covering-link S_n has two components, each of which is a trivial torus knot.

Before the proof, we give two known theorems.

Theorem 3.6 ([17], [2] p.73, 11.3 Theorem). *In the amalgamated product of groups G_i ($i = 1, 2$) with the amalgamated subgroup U , there is in each class of equivalent words one, and only one, element in a normal form $f = u z_1 z_2 \cdots z_t$. Here $u \in U$ and z_j ($j = 1, 2, \dots, t$) is a non-trivial right-handed coset representative of U in some $G_{i(j)}$, taken from an arbitrarily prefixed selection of right-handed coset representatives which contain the trivial coset representative, and $G_{i(j)} \neq G_{i(j+1)}$ for $j = 1, 2, \dots, t-1$.*

If the center of a classical knot group is non-trivial, then it is a torus knot (cf. [4]). There is a theorem concerning the center of a classical link group as follows.

Theorem 3.7 ([5]). *The statements listed below are equivalent.*

- (1) *The center of the group of a classical link L is non-trivial.*
- (2) *The link group is isomorphic to one of the groups of type (a), (b), or (c).*

$$(a) \underbrace{(Z * \cdots * Z)}_{m-1} \times \mathbb{Z},$$

$$(b) ((\underbrace{Z * \cdots * Z}_{m-2} \times \mathbb{Z}) *_{\mathbb{Z}} ((Z \times \mathbb{Z}) *_Z Z),$$

$$(c) \left(\underbrace{(Z * \cdots * Z)}_{m-2} \times \mathbb{Z} \right) *_\mathbb{Z} \left((\mathbb{Z} \times Z) *_\mathbb{Z} (Z *_Z Z) \right),$$

where m is the number of components of L , Z is an infinite cyclic group, and $\mathbb{Z} = \langle h \rangle$ is a “special” infinite cyclic group which is the center of the link group except when the link group is that of a Hopfian link of type (a). In case (b) the amalgamation concerning the last factor $Z = \langle q \rangle$ is given by $h = q^\alpha$ for an integer $\alpha > 1$. In case (c) the last factor $Z *_Z Z$ is the group of a torus (α, β) -knot.

Proof of Theorem 3.5. By Lemma 3.4, the link group G_n of S_n is computed as follows. Let $x_1, \dots, x_4$ be the generators. The relations concerning the vertical boundary braid $\sigma_1\sigma_3$ are $x_1 = x_2$ and $x_3 = x_4$. The other relations concerning the horizontal boundary braid Δ^{2n} are

$$\begin{aligned} x_1 &= (x_1x_2x_3x_4)^n x_1 (x_1x_2x_3x_4)^{-n}, \\ x_2 &= (x_1x_2x_3x_4)^n x_2 (x_1x_2x_3x_4)^{-n}, \\ x_3 &= (x_1x_2x_3x_4)^n x_3 (x_1x_2x_3x_4)^{-n}, \\ x_4 &= (x_1x_2x_3x_4)^n x_4 (x_1x_2x_3x_4)^{-n}. \end{aligned}$$

Putting $a = x_1 = x_2$ and $b = x_3 = x_4$, we have

$$G_n = \langle a, b \mid (a^2b^2)^n b = b(a^2b^2)^n, (a^2b^2)^n a = a(a^2b^2)^n \rangle.$$

Let Z_1 be the subgroup of G_1 generated by $\{a^2, b^2\}$, and Z_n be the subgroup of G_n generated by $h_n = (a^2b^2)^n$ for $n > 1$. Note that Z_n consists of central elements. We will show that Z_n is the center of G_n for $n \geq 1$. We can show as follows: Let N be a normal subgroup of G . If the center of G/N is trivial, then N is contained in the center of G . Hence it suffices to show that the center of the quotient group G_n/Z_n is trivial. If $n = 1$, then $G_1/Z_1 = \mathbb{Z}/2 * \mathbb{Z}/2$, whose center is trivial. If $n > 1$, then $G_n/Z_n = \langle a, b \mid (a^2b^2)^n = 1 \rangle$, which is an amalgamated product $\langle a \rangle *_U \langle b, x \mid x^n = 1 \rangle$, where $U = \langle a^2 \rangle = \langle xb^{-2} \rangle = \mathbb{Z}$ and the amalgamation is given by $a^2 = xb^{-2}$. Put $H_1 = \langle a \rangle$ and $H_2 = \langle b, x \mid x^n = 1 \rangle$. We can take $\{1, a\}$ as a set of right-handed coset representatives of U in H_1 . Take a set of right-handed coset representatives of U in H_2 , and denote it by C . Let h be a central element of $G_n/Z_n = H_1 *_U H_2$. By Theorem 3.6, h has a normal form $h = ua^\delta c_1 ac_2 \cdots ac_t a^\epsilon$, where $u \in U$ and $c_1, \dots, c_t \in C - \{1\}$ and $\delta, \epsilon \in \{0, 1\}$. Since $ah = ha$, we have $aua^\delta c_1 ac_2 \cdots ac_t a^\epsilon = ua^\delta c_1 ac_2 \cdots ac_t a^\epsilon a$. Since $ua = au$ in the amalgamated product $H_1 *_U H_2$, it follows that $ua^\delta ac_1 ac_2 \cdots ac_t a^\epsilon = ua^\delta c_1 ac_2 \cdots ac_t aa^\epsilon$, hence $ac_1 ac_2 \cdots ac_t = c_1 ac_2 \cdots ac_t a$ as elements in $H_1 *_U H_2$.

If $t > 0$, then $ac_1 ac_2 \cdots ac_t$ and $c_1 ac_2 \cdots ac_t a$ are in distinct normal forms, which is a contradiction. Hence $t = 0$ and we have $h = ua^\delta = a^k$ for an integer k . Since $hb = bh$, we have $a^k b = ba^k$. If $k = 1$, then we have $ab = ba$. In this case, if b is not in U , then we can take b as a non-trivial right-handed coset representative of U in H_2 . It follows that then ab and ba are in distinct normal forms, which is a contradiction. If $k = 2l + 1$ (resp. $k = 2l$) for a non-zero integer l , then we have $a^k b = uab$ and $ba^k = ca$ (resp. $a^k b = ub$ and $ba^k = c$), where in both cases $u = a^{2l} \in U$ and $c = b(xb^{-2})^l$. In these cases, if neither b nor c is in U and we can take b and c as distinct right-handed coset representatives of U in H_2 , then $a^k b$ and ba^k have distinct normal forms uab and ca (resp. ub and c), which is a contradiction. Then it follows that $k = 0$ and hence $h = 1$, i.e. the center of G_n/Z_n is trivial.

It remains to show that for non-zero integer l , neither b nor $c = b(xb^{-2})^l$ is in U and we can take b and c as distinct right-handed coset representatives of U in H_2 . The group $H_2 = \langle b, x \mid x^n = 1 \rangle$ is the free product of $\langle b \rangle$ and

$\langle x | x^n = 1 \rangle$. Hence by Theorem 3.6 again, every element of H_2 has a normal form $b^{\delta n_1} x^{m_1} b^{n_2} x^{m_2} \dots b^{n_t} x^{m_t}$, where $n_1, \dots, n_t \in \mathbb{Z} - \{0\}$, $m_1, \dots, m_t \in \{1, 2, \dots, n-1\}$, and $\delta, \epsilon \in \{0, 1\}$. Hence we can see that b has a normal form b , and $c = b(xb^{-2})^l$ has a normal form $b(xb^{-2})^{l_0}$ (resp. $b^3 x^{n-1} (b^2 x^{n-1})^{l_0-1}$) if $l > 0$ (resp. $l < 0$), where $l_0 = |l|$, a positive integer. On the other hand, an element of $U = \langle xb^{-2} \rangle$ in H_2 has a normal form $(xb^{-2})^{m+1}$ or $(b^2 x^{n-1})^m$, where m is a non-negative integer. Hence by the uniqueness of normal forms, we can see that neither b nor c is in U . Similarly, for a non-negative integer m , an element of Ub has a normal form $(xb^{-2})^m x b^{-1}$ or $(b^2 x^{n-1})^m b$, and an element of $Uc = Ub(xb^{-2})^l$ has a normal form $(xb^{-2})^m x b^{-1} (xb^{-2})^{l_0}$, $(xb^{-2})^m x b x^{n-1} (b^2 x^{n-1})^{l_0-1}$, $(b^2 x^{n-1})^m b (xb^{-2})^{l_0}$ or $(b^2 x^{n-1})^m b^3 x^{n-1} (b^2 x^{n-1})^{l_0-1}$, where $l_0 = |l|$, a positive integer. By the uniqueness of normal forms, we can see that $Ub \neq Uc$. Thus for non-zero integer l , neither b nor $c = b(xb^{-2})^l$ is in U and we can take b and c as distinct right-handed coset representatives of U in H_2 , and it follows that the center of G_n/Z_n is trivial.

Consider the abelianization map $\phi : G_n \rightarrow G_n/[G_n, G_n] = \mathbb{Z} \times \mathbb{Z}$ and put $\bar{a} = \phi(a)$ and $\bar{b} = \phi(b)$, which are the basis. Since $\phi(a^2) = \bar{a}^2$ and $\phi(b^2) = \bar{b}^2$, the center Z_1 is a rank-2 free abelian group. The image $\phi(h_n)$ is $\bar{a}^{2n} \bar{b}^{2n}$. Thus the center Z_n for $n > 1$ is an infinite cyclic group.

Next we will show that G_n is not a classical link group. By Theorem 3.7, we will show that G_n is neither of type (a), (b) nor (c) in Theorem 3.7. Since the torus-covering-link S_n consists of two components, it follows that m in Theorem 3.7 is two. We will first prove this for $n > 1$. Note that $\mathbb{Z}$ in Cases (b) and (c) in Theorem 3.7 is the center Z_n .

(Case (a)) If G_n is of type (a), then $G_n = Z \times \mathbb{Z}$, and it follows that G_n is commutative. On the other hand, there is a natural epimorphism

$$(3.2) \quad f : G_n \rightarrow \mathbb{Z}/2 * \mathbb{Z}/2 = \langle a' \rangle * \langle b' \rangle,$$

where $a' = f(a)$ and $b' = f(b)$, which are the basis. Since $\mathbb{Z}/2 * \mathbb{Z}/2$ is not commutative, G_n is not commutative. This is a contradiction.

(Case (b)) If G_n is of type (b), then $G_n = (Z \times Z_n) *_{Z_n} Z = (\langle k \rangle \times \langle h_n \rangle) *_{Z_n} \langle q \rangle$, where the amalgamation is given by $h_n = q^\alpha$ for an integer $\alpha > 1$. Put $h'_n = f(h_n)$ and $q' = f(q)$ for the natural epimorphism f of (3.2). Since $h_n = (a^2 b^2)^n$, we see that $h'_n = 1$. Since $h_n = q^\alpha$, it follows that $q'^\alpha = 1$. If $q' = 1$, then we have $f(G_n) = \langle f(k) \rangle$, which is generated by at most one generator. However, $f(G_n)$ is generated by two generators a' and b' . Hence q' is non-trivial. Consider the abelianization map $\phi' : f(G_n) \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2$, and put $\bar{a}' = \phi'(a')$ and $\bar{b}' = \phi'(b')$, which are the basis. Since $\alpha > 1$ and a non-trivial element of $f(G_n) = \mathbb{Z}/2 * \mathbb{Z}/2$ has order 2 or ∞ , we see that $\alpha = 2$, i.e. $q'^2 = 1$. Then it follows that $\phi'(q') = \bar{a}'$ or $\bar{b}'$. However, since $h_n = q^2$ and $\phi(h_n) = \bar{a}^{2n} \bar{b}^{2n}$, we have $\phi(q) = \bar{a}^n \bar{b}^n$. Then it follows that $\phi'(q') = 1$ or $\bar{a}' \bar{b}'$. This is a contradiction.

(Case (c)) If G_n is of type (c), then $G_n = (Z \times Z_n) *_{Z_n} (Z *_Z Z)$, where $Z *_Z Z$ is a classical knot group of a torus (α, β) -knot, i.e. $Z *_Z Z = \langle x, y | x^\alpha = y^\beta \rangle$ for coprime positive integers α and β . Then similarly to Case (b), we have the following argument. Since $h_n = (a^2 b^2)^n$, we see that $h'_n = 1$, where $h'_n = f(h_n)$ for the natural epimorphism f of (3.2). Since $h_n = x^\alpha = y^\beta$, it follows that $x'^\alpha = y'^\beta = 1$, where $x' = f(x)$ and $y' = f(y)$. If $x' = y' = 1$, then $f(\langle x, y | x^\alpha = y^\beta \rangle) = 1$ and it follows that $f(G_n)$ is generated by at most one generator. However, $f(G_n)$ is generated by two generators a' and b' . Hence we can assume that x' is non-trivial. Since any element of $f(G_n) = \mathbb{Z}/2 * \mathbb{Z}/2$ has order 2 or ∞ , it follows that $\alpha = 2$ and $\phi'(x') = \bar{a}'$ or $\bar{b}'$, where $\phi' : f(G_n) \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2$ is the abelianization map.

However, since $h_n = x^2$ and $\phi(h_n) = \bar{a}^{2n}\bar{b}^{2n}$, it follows that $\phi'(x')$ is 1 or $\bar{a}'\bar{b}'$. This is a contradiction.

In the case $n = 1$, since Z_1 is generated by two generators, if G_1 is a classical link group, then G_1 is of type (a). The rest of the proof is the same as in Case (a).

Now we will show that S_n and S_m are not equivalent for $n \neq m$. It suffices to show in the case when $n > 1$ and $m > 1$. Since the abelianization of G_n/Z_n is $\mathbb{Z} \times \mathbb{Z}/2n$, G_n is not isomorphic to G_m for $n \neq m$. $\square$

It is known that if the center of a μ -component 2-link group with $\mu > 1$ is non-trivial, then the center must be a torsion group (Corollary 2 of Chapter 3 in [9]). Hence we have a corollary.

Corollary 3.8. *The link groups of the 2-component torus-covering T^2 -links given in Theorem 3.5 are not 2-component 2-link groups.*

Proof. In the proof of Theorem 3.5, we have shown that the center of G_n , where $n > 0$, is non-trivial and torsion free. $\square$

We can consider the knot version of Theorem 3.5.

Theorem 3.9. *Let $\Gamma_{T,n}$ be the torus-covering-chart of degree 4 without black vertices and with boundary braids $\sigma_1\sigma_3$ and Δ^{2n+1} , where Δ is Garside's Δ and n is a positive integer. Let S_n be the torus-covering-knot associated with $\Gamma_{T,n}$. Then the knot group of S_n is not a classical knot group. Moreover, S_n and S_m are not equivalent for $n \neq m$.*

Remark. Note that S_0 is a trivial torus knot.

Proof. By Lemma 3.4, the knot group G_n of S_n is computed as follows. Let $x_1, \dots, x_4$ be the generators. Then the relations concerning the vertical boundary braid $\sigma_1\sigma_3$ are $x_1 = x_2$ and $x_3 = x_4$. The other relations concerning the horizontal boundary braid Δ^{2n+1} are

$$\begin{aligned} x_1 &= (x_1x_2x_3x_4)^n x_1x_2x_3x_4x_3^{-1}x_2^{-1}x_1^{-1}(x_1x_2x_3x_4)^{-n}, \\ x_2 &= (x_1x_2x_3x_4)^n x_1x_2x_3x_2^{-1}x_1^{-1}(x_1x_2x_3x_4)^{-n}, \\ x_3 &= (x_1x_2x_3x_4)^n x_1x_2x_1^{-1}(x_1x_2x_3x_4)^{-n}, \\ x_4 &= (x_1x_2x_3x_4)^n x_1(x_1x_2x_3x_4)^{-n}. \end{aligned}$$

Putting $a = x_1 = x_2$ and $b = x_3 = x_4$, we have

$$G_n = \langle a, b \mid b(a^2b^2)^n = (a^2b^2)^na, a(a^2b^2)^{n+1} = (a^2b^2)^{n+1}b \rangle.$$

Let Z_n be the subgroup of G_n generated by $h_n = (a^2b^2)^{2n+1}$, which is a central element. We show that Z_n is the center of G_n . Consider the quotient group $G'_n = G_n/Z_n$, which is $\langle a, b, x \mid x = a^2b^2, bx^n = x^na, ax^{n+1} = x^{n+1}b, x^{2n+1} = 1 \rangle$. By eliminating b by $b = x^na x^{-n}$, we have $G'_n = \langle a, x \mid x^{2n+1} = (a^2x^n)^2 = 1 \rangle$, which is an amalgamated product $\langle a \rangle *_U \langle x, y \mid x^{2n+1} = 1, y^2 = 1 \rangle$, where $U = \langle a^2 \rangle = \langle yx^{n+1} \rangle = \mathbb{Z}$ and the amalgamation is given by $a^2 = yx^{n+1}$. By an argument similar to the proof of Theorem 3.5, we can show that the center of G'_n is trivial. Thus Z_n is the center. Considering the abelianization map of G_n , we see that Z_n is an infinite cyclic group.

If G_n is a classical knot group, then it is a torus knot group. Let $G_{p,q}$ be the (p, q) -torus knot group isomorphic to G_n , where p and q are coprime positive integers. Let $Z_{p,q}$ be the center of $G_{p,q}$. The abelianization of G'_n is $\mathbb{Z}/4(2n+1)$. On the other hand, the abelianization of $G'_{p,q} = G_{p,q}/Z_{p,q}$ is $\mathbb{Z}/p \times \mathbb{Z}/q$. Since they are isomorphic, we have $pq = 4(2n+1)$. Since G'_n has an element of order

$2n + 1$, and the order of a non-trivial torsion element of $G'_{p,q}$ is a divisor of p or q , it follows that $2n + 1$ is a divisor of p or q . Hence we can determine coprime positive integers p and q to be $p = 4$ and $q = 2n + 1$.

For any element w of order 2 in $G'_{4,2n+1} = \mathbb{Z}/4 * \mathbb{Z}/(2n+1)$, w can be written as $w = w'^2$ for some element w' of order 4. Since y in G'_n is of order 2, there is an element y' with $y = y'^2$, and we have $G'_n = \langle a, x, y' \mid x^{2n+1} = 1, y'^4 = 1, a^2 x^n = y'^2 \rangle$. Let $N_{4,2n+1,v}$ be the normal subgroup of $G'_{4,2n+1}$ generated by an element v of order $2n+1$. Since the quotient group $G'_{4,2n+1}/N_{4,2n+1,v}$ does not depend on the choice of v , we will denote it by $G'_{4,2n+1}/N_{4,2n+1}$. Let N_n be the normal subgroup of G'_n generated by x . Since x has order $2n+1$, we see that G'_n/N_n is isomorphic to $G'_{4,2n+1}/N_{4,2n+1}$. We have $G'_n/N_n = \langle a, y' \mid y'^4 = 1, a^2 = y'^2 \rangle$, which is non-abelian. On the other hand, $G'_{4,2n+1}/N_{4,2n+1} = \mathbb{Z}/4$. This is a contradiction. Since the abelianization of G'_n is $\mathbb{Z}/4(2n+1)$, it follows that $S_n \neq S_m$ for $n \neq m$. $\square$

4. RIBBON TORUS-COVERING T^2 -LINKS

In this section we show Theorem 4.1.

Theorem 4.1. *The torus-covering T^2 -links given in Theorems 3.5 and 3.9 are ribbon but not symmetry-spun T^2 -links.*

Remark. Theorem 4.1 means that there are torus-covering T^2 -links which can be described by surface link charts without white vertices, but any torus-covering-chart describing which has white vertices.

Before the proof, we prove the following Proposition 4.3. First we will give the definition of *ribbon singularities* and certain conditions for boundary braids.

Let B^3 be a disjoint union of a finite number of handlebodies. The image of B^3 into $\mathbb{R}^4$ by an immersion ϕ is called a *3-ribbon* if the singularity set consists of *ribbon singularities*, i.e. the self-intersection of $\phi(B^3)$ consists of a finite number of mutually disjoint 2-disks, and for each 2-disk D_i , the preimage $\phi^{-1}(D_i)$ consists of a pair of 2-disks D'_i, D''_i such that $D'_i \cap D''_i = \emptyset$, $D'_i \subset \text{Int} B^3$ and $\partial D''_i = D'_i \cap \partial B^3$. An oriented surface link is ribbon if and only if it bounds a 3-ribbon (cf. [21]).

Let us consider a torus-covering-chart of degree nm . Let Γ_T be a torus-covering-chart of degree nm without black vertices and with boundary braids a and b . Let S be the torus-covering T^2 -link in $D^2 \times S^1 \times S^1$ associated with Γ_T , where D^2 is a 2-disk and $S^1 = [0, 1]/\sim$ with $0 \sim 1$. Let a_i (resp. b_i) be the i -th string of a (resp. b), where $i = 1, 2, \dots, nm$. A *cylinder* is $D_0^2 \times [0, 1]$, where D_0^2 is a 2-disk. Let us consider the following conditions.

- (R1) There are embedded cylinders $N_1(a), N_2(a), \dots, N_m(a)$ (resp. $N_1(b), N_2(b), \dots, N_m(b)$) in $D^2 \times [0, 1] \times \{0\}$ (resp. $D^2 \times \{0\} \times [0, 1]$) such that they are mutually disjoint in $D^2 \times [0, 1] \times \{0\}$ (resp. $D^2 \times \{0\} \times [0, 1]$) and $N_j(a) \cap a = a_{n(j-1)+1} \cup a_{n(j-1)+2} \cup \dots \cup a_{nj}$ (resp. $N_j(b) \cap b = b_{n(j-1)+1} \cup b_{n(j-1)+2} \cup \dots \cup b_{nj}$), where $j = 1, 2, \dots, m$.

By regarding each $N_j(a)$ (resp. $N_j(b)$) as a string, we have new classical m -braids. Let us call the new m -braid obtained from a (resp. b) the *tubular braid* associated with a (resp. b) and denote it by $R(a)$ (resp. $R(b)$).

- (R2) The associated tubular braid $R(a)$ is the trivial m -braid.

Let $R_1(a), R_2(a), \dots, R_m(a)$ be the n -braids associated with the vertical boundary braid a such that $R_j(a)$ corresponds to the j -th string of the trivial m -braid $R(a)$, where $j = 1, 2, \dots, m$. In other words, $R_j(a)$ is the n -braid $a \cap N_j(a)$ in the cylinder $N_j(a)$.

- (R3) The closed braid associated with $R_j(a)$ is a trivial knot, where $j = 1, 2, \dots, m$.




FIGURE 4.1. The boundary braids and the associated tubular braids

Let $N'\sigma_1$ be a $2n$ -braid defined as follows:

$$\begin{aligned}
 N'\sigma_1 &= \sigma_n(\sigma_{n-1}\sigma_{n-2}\cdots\sigma_1)(\sigma_{n+1}\sigma_{n+2}\cdots\sigma_{2n-1}) \\
 &\quad \cdot \sigma_n(\sigma_{n-1}\sigma_{n-2}\cdots\sigma_2)(\sigma_{n+1}\sigma_{n+2}\cdots\sigma_{2n-2}) \\
 &\quad \cdots \\
 &\quad \cdot \sigma_n(\sigma_{n-1})(\sigma_{n+1}) \\
 &\quad \cdot \sigma_n.
 \end{aligned}$$

Then for an m -braid β , let us define the nm -braid $N\beta$ as follows.


- (1) $N(\beta'\beta'') = N\beta' \cdot N\beta''$, for m -braids β' and β'' , and
- (2) $N\sigma_j = \iota_{n(j-1)}^{n(m-j-1)}(N'\sigma_1)$, where $j = 1, 2, \dots, m-1$.

Then under the conditions (R1), (R2) and (R3), the horizontal boundary braid b can be written uniquely as follows.

$$(4.1) \quad b = NR(b) \cdot \iota_0^{n(m-1)}(\beta_1) \cdot \iota_n^{n(m-2)}(\beta_2) \cdots \iota_{n(m-1)}^0(\beta_m),$$

where $\beta_1, \beta_2, \dots, \beta_m$ are n -braids.

Example 4.2. Let Γ_T be the torus-covering-chart of degree 4 without black vertices and with boundary braids $a = \sigma_1\sigma_3$ and $b = (\sigma_1\sigma_2\sigma_3)^4$. Note that b is equivalent to Δ^2 , where Δ is Garside's Δ (cf. Theorem 3.5). Then Γ_T satisfies the conditions (R1), (R2) and (R3). The associated tubular braids are 2-braids $R(a) = e$ (the trivial braid) and $R(b) = \sigma_1^2$. The 2-braids $R_1(a)$ and $R_2(a)$ are σ_1 , which are the braids associated with the vertical boundary braid a such that $R_1(a)$ (resp. $R_2(a)$) corresponds to the first (resp. second) string of the trivial 2-braid $R(a)$. See Fig. 4.1. Note that the closed braid associated with $R_j(a)$ ($j = 1, 2$) is a trivial knot. Since $R(b) = \sigma_1^2$, we have $NR(b) = N\sigma_1^2 = (\sigma_2\sigma_1\sigma_3\sigma_2)^2$, and the horizontal boundary braid b can be written uniquely as follows: $b = NR(b) \cdot \iota_0^2(\beta_1) \cdot \iota_2^0(\beta_2)$, where β_1 and β_2 are 2-braids σ_1^2 as in Fig. 4.2.

FIGURE 4.2. The boundary braid b

Proposition 4.3. *Let Γ_T be a torus-covering-chart of degree nm without black vertices and with boundary braids a and b such that the boundary braids a and b satisfy the conditions (R1), (R2) and (R3). Then the torus-covering T^2 -link associated with Γ_T is ribbon.*

Proof. Let S be the torus-covering T^2 -link associated with Γ_T , and we will use the notations in (R1), (R2), (R3) and (4.1). Let us prepare other notations. For a braid β , let $\text{cl}(\beta)$ be the closed braid associated with β . Let $E^3 = E^2 \times [0, 1]$ be a 3-ball which contain $D^2 \times S^1$, where E^2 is a 2-disk. The torus-covering-link S is embedded in $D^2 \times S^1 \times S^1$ in $E^3 \times S^1$, and $E^3 \times S^1$ is naturally embedded in $\mathbb{R}^4$.

(1) We consider the motion picture of S along S^1 of $E^3 \times S^1$, i.e. we consider S as $\cup_{t \in S^1} S_t$ in $E^3 \times S^1$, where $S_t = S \cap (E^3 \times \{t\})$. We can assume that $S^1 = [0, 2\nu + 2] / \sim$, where $0 \sim 2\nu + 2$. Divide S^1 into $2\nu + 2$ intervals $J_1^1, J_1^2, J_2^1, J_2^2, \dots, J_{\nu+1}^1, J_{\nu+1}^2$ with $J_k^1 = [2k - 2, 2k - 1]$ and $J_k^2 = [2k - 1, 2k]$, where $k = 1, 2, \dots, \nu + 1$. The associated tubular braid $R(b)$ is in $D^2 \times \{0\} \times [0, 2\nu + 2] \subset E^3 \times [0, 2\nu + 2]$. Let $R(b) = \sigma_{i_1}^{\epsilon_1} \cdot \sigma_{i_2}^{\epsilon_2} \cdots \sigma_{i_\nu}^{\epsilon_\nu}$ such that $R(b) \cap (D^2 \times \{0\} \times J_k^1) = \sigma_{i_k}^{\epsilon_k}$, where $i_k \in \{1, 2, \dots, m - 1\}$ and $\epsilon_k \in \{+1, -1\}$ for $k = 1, 2, \dots, \nu$. Then by (4.1) the horizontal boundary braid b can be written uniquely as follows. $b = N\sigma_{i_1}^{\epsilon_1} \cdot N\sigma_{i_2}^{\epsilon_2} \cdots N\sigma_{i_\nu}^{\epsilon_\nu} \cdot \iota_0^{n(m-1)}(\beta_1) \cdot \iota_n^{n(m-2)}(\beta_2) \cdots \iota_{n(m-1)}^0(\beta_m)$, where $\beta_1, \dots, \beta_m$ are n -braids.

Let $\{h_{k,u}\}_{u \in [0,1]}$ be an isotopy of D^2 associated with $N\sigma_{i_k}^{\epsilon_k}$ for $k = 1, 2, \dots, \nu$ (cf. Definition 3.1). Then let $\{f_{k,u}\}_{u \in [0,1]}$ be an isotopy of E^3 such that $f_u|_{D^2 \times S^1} = h_{k,u} \times \text{id}_{S^1}$ and $f_u|_{E^3 - D^2 \times S^1} = \text{id}_{E^3 - D^2 \times S^1}$, where $u \in [0, 1]$ and $k = 1, 2, \dots, \nu$. Similarly, let $\{h'_u\}_{u \in [0,1]}$ be an isotopy of D^2 associated with $\iota_0^{n(m-1)}(\beta_1) \cdot \iota_n^{n(m-2)}(\beta_2) \cdots \iota_{n(m-1)}^0(\beta_m)$. Then let $\{g_u\}_{u \in [0,1]}$ be an isotopy of E^3 such that $g_u|_{D^2 \times S^1} = h'_u \times \text{id}_{S^1}$ and $g_u|_{E^3 - D^2 \times S^1} = \text{id}_{E^3 - D^2 \times S^1}$, where $u \in [0, 1]$.

Then by definition of S , we can assume that the motion picture of S , i.e. $S = \cup_{t \in S^1} S_t$, where $S^1 = \cup_{k=1}^{\nu+1} (J_k^1 \cup J_k^2)$, is as follows. Identify $E^3 \times \{t\}$ with E^3 , where $t \in S^1$. Put $R_j^0 = R_j(a)$ for $j = 1, 2, \dots, m$ and $S_0 = \text{cl}(\iota_0^{n(m-1)}(R_1^0) \cdot \iota_n^{n(m-2)}(R_2^0) \cdots \iota_{n(m-1)}^0(R_m^0))$. Then

$$S_t = \begin{cases} f_{k,t-(2k-2)}(S_{2k-2}) & \text{if } t \in J_k^1 = [2k - 2, 2k - 1], \quad k = 1, 2, \dots, \nu, \\ g_{t-2\nu}(S_{2\nu}) & \text{if } t \in J_{\nu+1}^1 = [2\nu, 2\nu + 1], \\ S_{2k-1} & \text{if } t \in J_k^2 = [2k - 1, 2k], \quad k = 1, 2, \dots, \nu, \nu + 1. \end{cases}$$

Put

$$R_j^k = \begin{cases} R_j^k & \text{if } j \neq i_k - 1, i_k \\ R_{i_k}^k & \text{if } j = i_k - 1, \\ R_{i_k-1}^k & \text{if } j = i_k, \end{cases}$$

where $k = 0, 1, 2, \dots, \nu$ and $j = 1, 2, \dots, m$, and put

$$R^k = \iota_0^{n(m-1)}(R_1^k) \cdot \iota_n^{n(m-2)}(R_2^k) \cdots \iota_{n(m-1)}^0(R_m^k),$$

where $k = 0, 1, 2, \dots, \nu$. Then $R^{k-1} \cdot N\sigma_{i_k}^{\epsilon_k} = N\sigma_{i_k}^{\epsilon_k} \cdot R^k$ holds for $k = 1, 2, \dots, \nu$. The ambient isotopy $\{f_{k,u}\}$ is an ambient isotopy which deforms $\text{cl}(R^{k-1})$ to $\text{cl}(R^k)$. Let $R^{\nu+1} = R^0$. Note that

$$S_t = \text{cl}(R^k)$$

if $t \in J_k^2 = [2k-1, 2k]$, where $k = 1, 2, \dots, \nu, \nu+1$.

Since Γ_T has no black vertices, $ab = ba$ holds. Since $a = R^0$ and $a \cdot NR(b) = NR(b) \cdot R^\nu$, it follows that $R_j^\nu \beta_j = \beta_j R_j^0$ holds for $j = 1, 2, \dots, m$. The ambient isotopy $\{g_u\}$ is an ambient isotopy which deforms $S_{2\nu}$ to $S_{2\nu+1} = S_0$ as follows:

$$\begin{aligned} S_{2\nu} &= \text{cl}(R^\nu) \\ &= \text{cl}(\iota_0^{n(m-1)}(R_1^\nu) \cdot \iota_n^{n(m-2)}(R_2^\nu) \cdots \iota_{n(m-1)}^0(R_m^\nu)) \\ &= \text{cl}(\iota_0^{n(m-1)}(\beta_1 \cdot R_1^0 \cdot \beta_1^{-1}) \cdot \iota_n^{n(m-2)}(\beta_2 \cdot R_2^0 \cdot \beta_2^{-1}) \cdots \iota_{n(m-1)}^0(\beta_m \cdot R_m^0 \cdot \beta_m^{-1})) \\ &= \text{cl}(\iota_0^{n(m-1)}(\cdot R_1^0 \cdot \beta_1^{-1} \beta_1) \cdot \iota_n^{n(m-2)}(\cdot R_2^0 \cdot \beta_2^{-1} \beta_2) \cdots \iota_{n(m-1)}^0(\cdot R_m^0 \cdot \beta_m^{-1} \beta_m)) \\ &= \text{cl}(\iota_0^{n(m-1)}(R_1^0) \cdot \iota_n^{n(m-2)}(R_2^0) \cdots \iota_{n(m-1)}^0(R_m^0)) \\ &= \text{cl}(R^0) = S_0. \end{aligned}$$

(2) Next we will construct a 3-ribbon B such that $\partial B = S$.

Let $p : E^3 = E^2 \times [0, 1] \rightarrow E^2$ be the projection to the first factor. For two given subsets V, W of E^3 , we say V is over W with respect to p if for any point x in $p(V) \cap p(W)$, any pair of its preimages $(x, v) \in V$ and $(x, w) \in W$ satisfies that $v > w$, where $x \in E^2$ and $v, w \in [0, 1]$.

For $t \in S^1$, we identify $E^3 \times \{t\}$ with E^3 . Take mutually disjoint 2-disks $D_1^k, D_2^k, \dots, D_m^k$ in $E^3 \times \{2k\}$ such that $\partial D_j^k = \text{cl}(R_j^k)$ and D_j^k is over D_{j+1}^k with respect to the projection p for $j = 1, 2, \dots, m-1$ as follows, where $k = 0, 1, 2, \dots, \nu$. We take such 2-disks inductively for $k = 0, 1, 2, \dots, \nu$ as follows. First, since $\text{cl}(R_1^0), \text{cl}(R_2^0), \dots, \text{cl}(R_m^0)$ are trivial knots by (R3), we can take such 2-disks $D_1^0, D_2^0, \dots, D_m^0$. Then define $D_j^1 = f_{1,1}(D_j^0)$ if $j \neq i_1 - 1, i_1$, where $j = 1, 2, \dots, m$, and $D_{i_1-1}^1 = f_{1,1}(D_{i_1}^0)$. And then take a 2-disk $D_{i_1}^1$ with $\partial D_{i_1}^1 = \text{cl}(R_{i_1}^1)$ which satisfies the conditions that (i) $D_{i_1}^1 \cap D_j^1 = \emptyset$ if $j \neq i_1$, where $j = 1, 2, \dots, m-1$, and (ii) D_j^1 is over D_{j+1}^1 with respect to p for $j = 1, 2, \dots, m-1$. Then define $D_j^2 = f_{2,1}(D_j^1)$ if $j \neq i_2 - 1, i_2$, where $j = 1, 2, \dots, m$, and $D_{i_2-1}^2 = f_{2,1}(D_{i_2}^1)$. And then take a 2-disk $D_{i_2}^2$ with $\partial D_{i_2}^2 = \text{cl}(R_{i_2}^2)$ which satisfies the conditions that (i) $D_{i_2}^2 \cap D_j^2 = \emptyset$ if $j \neq i_2$, where $j = 1, 2, \dots, m$, and (ii) D_j^2 is over D_{j+1}^2 with respect to p for $j = 1, 2, \dots, m-1$. Repeating this process, we have 2-disks $D_1^k, D_2^k, \dots, D_m^k$ for $k = 0, 1, 2, \dots, \nu$. Note that we have

$$D_j^k = \begin{cases} f_{k,1}(D_j^{k-1}) & \text{if } j \neq i_k - 1, i_k \\ f_{k,1}(D_{i_k}^{k-1}) & \text{if } j = i_k - 1, \end{cases}$$

where $j = 1, 2, \dots, m$ and $k = 1, 2, \dots, \nu$.

Let us denote $f_{k,1}(D_{i_k}^{k-1})$ by $D_{i_k}^k$. Take an ambient isotopy $\{\phi_{k,u}\}_{u \in [0,1]}$ which carries $D_{i_k}^{k-1}$ to $D_{i_k}^k$, where $k = 1, 2, \dots, \nu$, and let $\{\psi_u\}_{u \in [0,1]}$ be an ambient isotopy which carries $g_1(D_1^\nu \cup D_2^\nu \cup \dots \cup D_m^\nu)$ to $D_1^0 \cup D_2^0 \cup \dots \cup D_m^0$. Since D_j^ν (resp. D_j^0) is over D_{j+1}^ν (resp. D_{j+1}^0) with respect to p for $j = 1, 2, \dots, m-1$, $\{\psi_u\}$ exists.

For $k = 1, 2, \dots, \nu + 1$ and $j = 1, 2, \dots, m$, define $D_j^k(t)$ for $t \in J_k^1$ and $t \in J_k^2$ as follows.

$$D_j^k(t) = \begin{cases} f_{k, t-(2k-2)}(D_j^{k-1}) & \text{if } t \in J_k^1 = [2k-2, 2k-1], \quad k = 1, 2, \dots, \nu, \\ D_j^k & \text{if } t \in J_k^2 = [2k-1, 2k], \quad j \neq i_k, \quad k = 1, 2, \dots, \nu, \\ \phi_{k, t-(2k-1)}(D_{i_k}^k) & \text{if } t \in J_k^2 = [2k-1, 2k], \quad j = i_k, \quad k = 1, 2, \dots, \nu, \\ g_{t-2\nu}(D_j^\nu) & \text{if } t \in J_{\nu+1}^1 = [2\nu, 2\nu+1], \quad k = \nu+1, \\ \psi_{t-(2\nu+1)}(D_j^\nu) & \text{if } t \in J_{\nu+1}^2 = [2\nu+1, 2\nu+2], \quad k = \nu+1, \end{cases}$$

where $j = 1, 2, \dots, m$.

Then put

$$B_t = D_1^k(t) \cup \dots \cup D_m^k(t),$$

where $t \in \cup_{k=1}^{\nu+1} (J_k^1 \cup J_k^2)$, and define

$$B = \cup_{t \in S^1} B_t = \cup_{k=1}^{\nu+1} \cup_{t \in J_k^1 \cup J_k^2} B_t.$$

We can assume that $D_{i_k}^k(t)$ is disjoint with $\cup_{j \neq i_k-1, i_k} D_j^k(t)$, where $t \in J_k^2$ and $k = 1, 2, \dots, \nu$. By the construction, $(\cup_{k=1}^{\nu+1} \cup_{t \in J_k^1} B_t) \cup (\cup_{t \in J_{\nu+1}^2} B_t)$ has no singularities, and for $k = 1, 2, \dots, \nu$, $\cup_{t \in J_k^2} B_t$ has n singular 2-disks whose singularities are ribbon. Hence B is a 3-ribbon such that $\partial B = S$. $\square$

Proof of Theorem 4.1. By Proposition 4.3, the torus-covering T^2 -links given in Theorems 3.5 and 3.9 are ribbon. $\square$

5. TRIPLE POINT NUMBERS

The triple point number of a surface link F is the minimum number of triple points in a surface diagram of F , for all the surface diagrams. Symmetry-spun T^2 -links are pseudo-ribbon, and by Theorem 4.1 the torus-covering T^2 -links given in Theorems 3.5 and 3.9 are ribbon. Both of them have the triple point number zero. In this section we give an example of a torus-covering T^2 -knot whose triple point number is positive.

Theorem 5.1. *Let Γ_T be the torus-covering-chart of degree 4 without black vertices and with boundary braids $\sigma_1\sigma_2^3\sigma_3$ and $(\sigma_1\sigma_2\sigma_3)^4$. Then the triple point number of the torus-covering-knot S associated with Γ_T is positive.*

We tri-color the surface diagram associated with Γ_T and show that the quandle cocycle invariant associated with Mochizuki's 3-cocycle does not have an integer value. Then any surface diagram of S has at least four triple points (cf. [18]). We use the following facts. Let F be an oriented surface link.

Let $\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a generic projection. In the surface diagram $D = \pi(F)$, there are two intersecting sheets along each double point curve, one of which is higher than the other with respect to π . They are called the *over sheet* and the *under sheet* along the double point curve, respectively. In order to indicate crossing information of the surface diagram, we break the under sheet into two pieces missing the over sheet. This can be extended around a triple point. Around a triple point, the sheets are called the *top sheet*, the *middle sheet*, and the *bottom sheet* from the higher one. Then the surface diagram is presented by a disjoint union of compact surfaces which are called *broken sheets*. We denote by $B(D)$ the set of broken sheets of D .

A set X with a binary operation $*$: $X \times X \rightarrow X$ is called a *quandle* if it satisfies the following conditions:

- (i) for any $a \in X$, $a * a = a$,
- (ii) for any $a, b \in X$, there exists a unique $c \in X$ such that $a = c * b$, and

(iii) for any $a, b, c \in X$, $(a * b) * c = (a * c) * (b * c)$ (cf. [11]).

The *dihedral quandle* of order 3, R_3 , is the set $\{0, 1, 2\}$ with the binary operation $a * b = 2b - a \pmod{3}$. A *tri-coloring* for a surface diagram D is a map $C : B(D) \rightarrow R_3$ such that $C(H_1) * C(H_2) = C(H'_1)$ along each double point curve of D , where H_2 is the over sheet and H_1 (resp. H'_1) is the under sheet such that the normal vector of H_2 points from (resp. toward) it. The image by C is called the *color*.

At a triple point of D , there exist broken sheets $J_1, J_2, J_3 \in B(D)$ uniquely such that J_1 is the bottom sheet, J_2 is the middle sheet, J_3 is the top sheet and the normal vector of J_2 (resp. J_3) points from J_1 (resp. J_2). The *color of the triple point* is the triplet $(C(J_1), C(J_2), C(J_3)) \in R_3 \times R_3 \times R_3$. The *sign* of the triple point is *positive* or $+1$ (resp. *negative* or -1) if the triplet of the normal vectors of J_1, J_2, J_3 is right-handed (resp. left-handed).

If D has a corresponding chart, this corresponds to the following (cf. Proposition 4.43 (3) in [8]). The color of the white vertex representing $\sigma_i \sigma_j \sigma_i \rightarrow \sigma_j \sigma_i \sigma_j$ ($|i - j| = 1$) is (a, b, c) , where a, b and c are the colors of the broken sheets of D connected with the starting points of the i' -th, $(i' + 1)$ -th, and $(i' + 2)$ -th strings of $\sigma_i \sigma_j \sigma_i$, where $i' = \min\{i, j\}$. The white vertex is positive (resp. negative) if $j > i$ (resp. $i > j$), i.e. if there is exactly one edge with the largest (resp. smallest) label oriented toward the white vertex.





Mochizuki's 3-cocycle is a map $\theta : R_3 \times R_3 \times R_3 \rightarrow \mathbb{Z}_3 = \langle t \mid t^3 = 1 \rangle$ such that $\theta(x, y, z) = t^{(x-y)(y-z)z(x+z)} \in \mathbb{Z}_3$.

Let $\tau_1, \dots, \tau_s$ be all the triple points of D with the sign $\epsilon_i \in \{+1, -1\}$ and the color $(a_i, b_i, c_i) \in R_3 \times R_3 \times R_3$ for each τ_i . Put $W_\theta(\tau_i; C) = \theta(a_i, b_i, c_i)^{\epsilon_i} \in \mathbb{Z}_3$ for each τ_i , and $W_\theta(C) = \prod_{i=1}^s W_\theta(\tau_i; C) \in \mathbb{Z}_3$ for the tri-coloring C . We call $W_\theta(C)$ the *Boltzman weight*. Since the set of broken sheets of D is finite, so is the set of tri-colorings for D . Let $C_1, \dots, C_n$ be all the tri-colorings, and define $\Phi_\theta(D)$ by $\Phi_\theta(D) = \sum_{j=1}^n W_\theta(C_j) \in \mathbb{Z}\langle t \mid t^3 = 1 \rangle$. Since θ satisfies certain conditions, $\Phi_\theta(D)$ does not depend on the choice of the surface diagram D of F . We call $\Phi_\theta(D)$ the *quandle cocycle invariant* of F associated with Mochizuki's 3-cocycle, and use the notation $\Phi_\theta(F)$.

Remark. For the quandle cocycle invariants, there is a general theory (cf. [6, 7]).

Proof of Theorem 5.1. Tri-color the surface diagram associated with Γ_T . Then the tri-coloring for the diagrams of the boundary braids $\sigma_1 \sigma_2^3 \sigma_3$ and $(\sigma_1 \sigma_2 \sigma_3)^4$ is as in Figs 5.1 and 5.2, where $\{a, b, c\} = \{0, 1, 2\}$, $\{0\}$, $\{1\}$ or $\{2\}$, which are all the tri-colorings. Let us denote by C the tri-coloring described by $\{a, b, c\}$. The part of the torus-covering-chart without black vertices and with boundary braids σ_1 and $(\sigma_1 \sigma_2 \sigma_3)^4$ has four white vertices. Denote these by $\tau_{11}^1, \dots, \tau_{14}^1$ from left to right as in Fig. 5.3. The color of each white vertex is obtained from reading the colors along the dotted path in Fig. 5.2, and the sign is obtained from the labels and the orientations of the edges around the white vertex. Similarly, we have white vertices τ_{ij}^2 ($i = 2, 3, 4, j = 1, 2, 3, 4$) and $\tau_{51}^3, \dots, \tau_{54}^3$. Fig. 5.4 shows the white vertices $\tau_{i1}^2, \dots, \tau_{i4}^2$ and the colors when $i = 2$. The matrix which describes the sign and the color of each white vertex is as follows, where the (i, j) -element describes those of τ_{ij}^* :

$$\begin{pmatrix} +(a, a, c) & +(b, b, c) & -(c, a, a) & -(c, a, a) \\ -(a, a, c) & +(c, b, c) & +(c, a, a) & -(b, a, c) \\ -(a, c, b) & +(b, a, c) & +(a, b, a) & -(b, c, b) \\ -(a, b, a) & +(a, c, c) & +(b, c, a) & -(b, b, a) \\ -(a, c, c) & -(a, c, c) & +(c, c, a) & +(b, b, a) \end{pmatrix}.$$

FIGURE 5.1. The tri-coloring for $\sigma_1\sigma_2^3\sigma_3$ FIGURE 5.2. The tri-coloring for $(\sigma_1\sigma_2\sigma_3)^4$ FIGURE 5.3. White vertices $\tau_{11}^1, \dots, \tau_{14}^1$ FIGURE 5.4. White vertices $\tau_{i1}^2, \dots, \tau_{i4}^2$ and the colors when $i = 2$

Then the Boltzman weight $W_\theta(C)$ for the Mochizuki's 3-cocycle θ and the tri-coloring C is $W_\theta(C) = \theta(c, b, c) \cdot \theta(a, c, b)^{-1} \cdot \theta(b, c, b)^{-1} \cdot \theta(b, c, a)$, and we have

$$W_\theta(C) = \begin{cases} t^2 & \text{if } \{a, b, c\} = \{0, 1, 2\} \\ 1 & \text{if } \{a, b, c\} = \{0\}, \{1\}, \{2\}. \end{cases}$$

Hence $\Phi_\theta(S)$, the quandle cocycle invariant for S associated with Mochizuki's 3-cocycle, is $\Phi_\theta(S) = 3 + 6t^2 \notin \mathbb{Z}$ in $\mathbb{Z}[t]/\langle t^3 = 1 \rangle$. $\square$

An oriented surface link F is *(-)-amphicheiral* if $F = -F^*$, where $-F^*$ is the orientation-reversed mirror image of F . Symmetry-spun T^2 -links are *(-)-amphicheiral*. Similarly to the proof of Theorem 5.1, using $-\Gamma_T^*$, we can see that $\Phi_\theta(-S^*) = 3 + 6t \neq \Phi_\theta(S)$. Thus we have a corollary.

Corollary 5.2. *The torus-covering T^2 -knot of Theorem 5.1 is not *(-)-amphicheiral*.*

REFERENCES

- [1] J. Birman, A. Lubotzky, and J. McCarthy, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. **50** (1983), 1107–1120.
- [2] O. Bogopolski, *Introduction to Group Theory*, European Mathematical Society, 2008.
- [3] J. Boyle, *The turned torus knot in S^4* , J. Knot Theory Ramifications **2** (1993), 239–249.
- [4] G. Burde and H. Zieschang, *Eine Kennzeichnung der Torusknoten*, Math. Ann. **169** (1966), 169–176.
- [5] G. Burde and K. Murasugi, *Links and Seifert fiber spaces*, Duke Math. J. **37** (1970), 89–93.
- [6] J. S. Carter, D. Jelsovsky, S. Kamada, L. Langford and M. Saito, *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, Trans. Amer. Math. Soc. **355** (2003), 3947–3989.
- [7] J. S. Carter, D. Jelsovsky, S. Kamada, and M. Saito, *Computations of quandle cocycle invariants of knotted curves and surfaces*, Adv. in Math. **157** (2001), 36–94.
- [8] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, Mathematical Surveys and Monographs 55, Amer. Math. Soc., 1998.
- [9] J. Hillman, *2-Knots and their Groups*, Australian Mathematical Society Lecture Series. 5, Cambridge University Press, 1989.
- [10] Z. Iwase, *Dehn-surgery along a torus T^2 -knot*, Pacific J. Math. **133** (1988), 289–299.
- [11] D. Joyce, *A classical invariant of knots, the knot quandle*, J. Pure Appl. Algebra **23** (1982), 37–65.
- [12] S. Kamada, *A characterization of groups of closed orientable surfaces in 4-space*, Topology **33** (1994), 113–122.
- [13] S. Kamada, *An observation of surface braids via chart description*, J. Knot Theory Ramifications **4** (1996), 517–529.
- [14] S. Kamada, *Braid and Knot Theory in Dimension Four*, Math. Surveys and Monographs 95, Amer. Math. Soc., 2002.
- [15] A. Kawauchi, *On pseudo-ribbon surface-links*, J. Knot Theory Ramifications **11** No.7 (2002), 1043–1062.
- [16] R. A. Litherland, *The second homology of the group of a knotted surface*, Quart. J. Math. Oxford (2) **32** (1981), 425–434.
- [17] H. Neumann, *Generalized free products with amalgamated subgroups*, Amer. J. Math., **71** (1949), 491–540.
- [18] S. Satoh and A. Shima, *The 2-twist-spun trefoil has the triple point number four*, Trans. Amer. Math. Soc. **356** (2004), 1007–1024.
- [19] A. Shima, *Knotted Klein bottles with only double points*, Osaka J. Math. **40** (2003), 779–799.
- [20] M. Teragaito, *Symmetry-spun tori in the four sphere*, Knots 90, 163–171.
- [21] T. Yanagawa, *On ribbon 2-knots*, Osaka J. Math. **6** (1969), 447–464.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO
E-mail address: inasa@ms.u-tokyo.ac.jp